

**SELF-INTERSECTIONS FOR THE SURFACE
DIFFUSION AND THE VOLUME-PRESERVING
MEAN CURVATURE FLOW**

UWE F. MAYER AND GIERI SIMONETT

Department of Mathematics, Vanderbilt University, Nashville, TN 37240

(Submitted by: Herbert Amann)

Abstract. We prove that the surface-diffusion flow and the volume-preserving mean curvature flow can drive embedded hypersurfaces to self-intersections.

1. Introduction. In this paper we consider two geometric evolution laws: the surface-diffusion flow and the volume-preserving mean curvature flow. We prove that embedded hypersurfaces can be driven to a self-intersection in finite time. This situation is in strict contrast to the behavior of hypersurfaces under the mean curvature flow, where the maximum principle prevents self-intersections.

The surface-diffusion flow is a geometric evolution law in which the normal velocity of a moving hypersurface equals the Laplace-Beltrami of the mean curvature. More precisely we assume in the following that Γ_0 is a closed, embedded hypersurface in \mathbb{R}^n . Then the surface-diffusion flow is governed by the law

$$V(t) = \Delta_{\Gamma(t)} H_{\Gamma(t)}, \quad \Gamma(0) = \Gamma_0. \quad (1.1)$$

Here $\Gamma = \{\Gamma(t) : t \geq 0\}$ is a family of smooth, immersed, orientable hypersurfaces, $V(t)$ denotes the velocity of Γ in the normal direction at time t , while $\Delta_{\Gamma(t)}$ and $H_{\Gamma(t)}$ stand for the Laplace-Beltrami operator and the mean curvature of $\Gamma(t)$, respectively. The volume-preserving mean curvature flow is governed by the law

$$V(t) = -(H(t) - \overline{H}(t)), \quad \Gamma(0) = \Gamma_0, \quad (1.2)$$

Accepted for publication May 1999.

AMS Subject Classifications: 35R35, 35K55, 58G11, 58F39.

where $\overline{H}(t) := |\Gamma(t)|^{-1} \int_{\Gamma(t)} H(t) d\sigma$ denotes the average of the mean curvature.

The evolution laws (1.1) and (1.2) do not depend on the local choice of the orientation. However, if $\Gamma(t)$ is embedded and encloses a region $\Omega(t)$ we always choose the outer normal, so that $V(t)$ is positive if $\Omega(t)$ grows, and so that $H_{\Gamma(t)}$ is positive if $\Gamma(t)$ is convex with respect to $\Omega(t)$.

The surface-diffusion flow (1.1) was first proposed by Mullins [20] to model the dynamics for the motion of the surface of a crystal when all mass transport is by curvature-driven diffusion along the surface. It has also been examined in a more general mathematical and physical context by Davi and Gurtin [10], and by Cahn and Taylor [7]. The surface-diffusion flow has recently attracted the attention of various researchers; see [3, 4, 6, 8, 11, 12, 13, 16, 19, 21]. We refer to [5, 14, 15, 17] for work related to the volume-preserving mean curvature flow.

The surface-diffusion flow and the volume-preserving mean curvature flow evolve hypersurfaces in such a way that the surface area decreases. Moreover, if Γ is embedded then both flows preserve the volume of the region $\Omega(t)$ enclosed by $\Gamma(t)$; see for instance [12, 14]. The results herein show that both flows can force $\Gamma(t)$ to lose embeddedness in order to decrease surface area.

Theorem 1. *Let $0 < \beta < 1$. There exist a closed embedded hypersurface $\Sigma_0 \in C^{2+\beta}$, a constant $T_0 > 0$, numbers $t_0, t_1 \in (0, T_0]$ with $t_0 < t_1$, and a $C^{2+\beta}$ -neighborhood U_0 of Σ_0 such that the surface-diffusion flow (1.1) has a unique classical solution $\Gamma = \{\Gamma(t) : t \in [0, T_0]\}$ for all $\Gamma_0 \in U_0$, and such that $\Gamma(t)$ ceases to be embedded for every $t \in (t_0, t_1)$ and every $\Gamma_0 \in U_0$. Each hypersurface $\Gamma(t)$ is of class C^∞ for $t \in (0, T_0]$ and smooth in $t \in (0, T_0)$.*

It was conjectured in [11] and later proved in [16] that the surface-diffusion flow can drive a dumbbell curve of an appropriate shape to a self-intersection. Theorem 1 extends this result considerably: we can handle nonsymmetric hypersurfaces in any dimension, whereas the method of [16] seems restricted to (symmetric) curves. It should be noted that the neighborhood U_0 of Theorem 1 also contains C^∞ -hypersurfaces that will be driven to a self-intersection in finite time. Our approach relies on results and techniques in [12].

Theorem 2. *Let $0 < \beta < 1$. There exist a closed embedded hypersurface $\Sigma_0 \in C^{1+\beta}$, a constant $T_0 > 0$, numbers $t_0, t_1 \in (0, T_0]$ with $t_0 < t_1$, and a*

$C^{1+\beta}$ -neighborhood U_0 of Σ_0 such that the volume-preserving mean curvature flow (1.2) has a unique classical solution $\Gamma = \{\Gamma(t) : t \in [0, T_0]\}$ for all $\Gamma_0 \in U_0$, and such that $\Gamma(t)$ ceases to be embedded for every $t \in (t_0, t_1)$ and every $\Gamma_0 \in U_0$. Each hypersurface $\Gamma(t)$ is of class C^∞ for $t \in (0, T_0]$ and smooth in $t \in (0, T_0)$.

To the best of our knowledge, Theorem 2 provides the first rigorous proof for the occurrence of self-intersections for the volume-preserving mean curvature flow. In particular, we give a proof for an example proposed by Gage [15] who considered a curve similar to our Figure 3.

2. The surface-diffusion flow. In this section we prove Theorem 1. We first introduce some notations. Given an open set $U \subset \mathbb{R}^n$, let $h^s(U)$ denote the little Hölder spaces of order $s > 0$, that is, the closure of $BUC^\infty(U)$ in $BUC^s(U)$, the latter space being the Banach space of all bounded and uniformly Hölder-continuous functions of order s . If Σ is a (sufficiently) smooth submanifold of \mathbb{R}^n then the spaces $h^s(\Sigma)$ are defined by means of a smooth atlas for Σ . It is known that $BUC^t(\Sigma)$ is continuously embedded in $h^s(\Sigma)$ whenever $t > s$. In the following, we assume that Σ is a smooth, compact, closed, immersed, oriented hypersurface in \mathbb{R}^n . Let ν be the unit normal field on Σ commensurable with the chosen orientation. Then we can find $a > 0$ and an open covering $\{U_l : l = 1, \dots, m\}$ of Σ such that

$$X_l : U_l \times (-a, a) \rightarrow \mathbb{R}^n, \quad X_l(s, r) := s + r\nu(s)$$

is a smooth diffeomorphism onto its image $\mathcal{R}_l := \text{im}(X_l)$; that is,

$$X_l \in \text{Diff}^\infty(U_l \times (-a, a), \mathcal{R}_l), \quad 1 \leq l \leq m.$$

This can be done by selecting the open sets $U_l \subset \Sigma$ in such a way that they are embedded in \mathbb{R}^n instead of only immersed, and then taking $a > 0$ sufficiently small so that each of the U_l has a tubular neighborhood of radius a . It follows that $\mathcal{R} := \cup \mathcal{R}_l$ consists of those points in \mathbb{R}^n with distance less than a to Σ . Let $\beta \in (0, 1)$ be fixed. Then we choose numbers $\alpha, \beta_0 \in (0, 1)$ with $\alpha < \beta_0 < \beta$. Let

$$W := \{\rho \in h^{2+\beta_0}(\Sigma) : \|\rho\|_\infty < a\}. \tag{2.1}$$

Given any $\rho \in W$ we obtain a compact, oriented, immersed manifold Γ_ρ of class $h^{2+\beta_0}$ by means of the following construction:

$$\Gamma_\rho := \bigcup_{l=1}^m \text{Im} (X_l : U_l \rightarrow \mathbb{R}^n, [s \mapsto X_l(s, \rho(s))]).$$

Thus Γ_ρ is a graph in normal direction over Σ and ρ is the signed distance between Σ and Γ_ρ . On the other hand, every compact, immersed, oriented manifold Γ that is a smooth graph over Σ and that is contained in \mathcal{R} can be obtained in this way. For convenience we introduce the mapping

$$\theta_\rho : \Sigma \rightarrow \Gamma_\rho, \quad \theta_\rho(s) := X_l(s, \rho(s)) \text{ for } s \in U_l, \quad \rho \in W.$$

It follows that θ_ρ is a well-defined global $(2+\beta_0)$ -diffeomorphism from Σ onto Γ_ρ . The surface-diffusion flow (1.1) can now be expressed as an evolution equation for the distance function ρ over the fixed reference manifold Σ ,

$$\partial_t \rho = G(\rho), \quad \rho(0) = \rho_0. \quad (2.2)$$

Here $G(\rho) := L_\rho \theta_\rho^*(\Delta_{\Gamma_\rho} H_{\Gamma_\rho})$ for $\rho \in h^{4+\alpha}(\Sigma) \cap W$, while Δ_{Γ_ρ} and H_{Γ_ρ} are the Laplace-Beltrami operator and the mean curvature of Γ_ρ , respectively, and $L(\rho)$ is a factor that comes in by calculating the normal velocity in terms of ρ ; see [12] for more details. We are now ready to state the following existence result for solutions of (2.2).

Proposition 2.1.

(a) *Let $\sigma \in W$ be given. Then there exist a positive constant $T_0 > 0$ and a neighborhood $W_0 \subset W$ of σ in $h^{2+\beta_0}(\Sigma)$ such that (2.2) has a unique solution,*

$$\rho(\cdot, \rho_0) \in C([0, T_0], W) \cap C^\infty((0, T_0) \times \Sigma) \text{ for every } \rho_0 \in W_0.$$

(b) *The map $[(t, \rho_0) \mapsto \rho(t, \rho_0)]$ defines a smooth local semiflow on W_0 .*

(c) *$\rho(\cdot, \rho_0) \in C([0, T_0], h^{4+\alpha}(\Sigma)) \cap C^1([0, T_0], h^\alpha(\Sigma))$ for every $\rho_0 \in h^{4+\alpha}(\Sigma) \cap W_0$.*

Proof. (a) and (b) follow from [12, Theorem 2.2]. Moreover, [12, Lemma 2.1] shows that the mapping $[\rho \mapsto G(\rho)] : h^{4+\alpha}(\Sigma) \cap W \rightarrow h^\alpha(\Sigma)$ is smooth and that the derivative is given by $G'(\rho) = P(\rho) + B(\rho)$, where

$$P(\rho) \in L(h^{4+\alpha}(\Sigma), h^\alpha(\Sigma)), \quad B(\rho) \in L(h^{2+\alpha}(\Sigma), h^\alpha(\Sigma)), \quad \rho \in h^{4+\alpha}(\Sigma) \cap W.$$

In the following we fix $\rho \in h^{4+\alpha}(\Sigma) \cap W$. Lemma 2.1 in [12] also shows that $P(\rho)$ generates a strongly continuous analytic semigroup on $h^\alpha(\Sigma)$. A well-known perturbation result, see [1, Theorem I.1.3.1], then implies $G'(\rho) \in L(h^{4+\alpha}(\Sigma), h^\alpha(\Sigma))$ also generates a strongly continuous analytic semigroup on $h^\alpha(\Sigma)$. It is known (see [1, Vol II], for instance) that the little Hölder

spaces are stable under the continuous interpolation method [1, 2, 9, 18]. Therefore, the spaces $(h^{4+\alpha}(\Sigma), h^\alpha(\Sigma))$ form a pair of maximal regularity for $G'(\rho)$; see [1, Theorem III.3.4.1] or [2, 9, 18]. Part (c) follows now from maximal regularity results, for instance [2, Theorem 2.7]. \square

In order to provide a proof of Theorem 1 we now choose Σ to be any smooth, compact, closed, immersed, orientable hypersurface in \mathbb{R}^n such that its image contains the flat $(n - 1)$ -dimensional disk $U := \{(s, 0) \in \mathbb{R}^{n-1} \times \mathbb{R} : |s| \leq 1\}$ twice, and with opposite orientations. To be precise, let $i : \Sigma \rightarrow \mathbb{R}^n$ be the immersion under consideration; then we ask that

$$i^{-1}(U) = U^+ \cup U^-$$

with $U^+ \cap U^- = \emptyset$ and both U^+ and U^- are flat $(n - 1)$ -dimensional disks of radius 1. Additionally we ask that $\Sigma \setminus (U^+ \cup U^-)$ be embedded in \mathbb{R}^n . Identifying U^+ for the moment with its image U we ask that the normal on U^+ points upwards; that is, $\nu(\cdot)|_{U^+} = e_n$, the n^{th} basis vector of \mathbb{R}^n . It follows that $\nu(\cdot)|_{U^-} = -e_n$.




Figure 1 This is a possible choice of Σ , cut in halves.

Let W be as in (2.1), and let $\sigma \in h^{4+\alpha} \cap W$ have the following local symmetry: $\sigma(-s) = \sigma(s)$ for every $s \in U^\pm$. This implies $\partial_j \sigma(0) = 0$ for $1 \leq j \leq n - 1$. Observe that $\theta_\sigma(s) = (s, \pm \sigma(s))$ (these are coordinates in \mathbb{R}^n) for $s \in U^\pm$ and that $\theta_\sigma : U^\pm \rightarrow \theta_\sigma(U^\pm)$ is an $h^{4+\alpha}$ -diffeomorphism. It is straightforward to compute

$$G(\sigma)|_{U^\pm} := L(\rho)\theta_\sigma^*(\Delta_{\Gamma_\sigma} H_{\Gamma_\sigma})|_{U^\pm}$$

in local coordinates, yielding

$$\begin{aligned}
 (n-1)G(\sigma)|_{U^\pm}(0) &= -\Delta_{n-1}^2\sigma(0) + \sum_{j,k=1}^{n-1} (\partial_j\partial_k\sigma(0))^2\Delta_{n-1}\sigma(0) \\
 &\quad + 2 \sum_{j,k,l=1}^{n-1} \partial_j\partial_k\sigma(0)\partial_j\partial_l\sigma(0)\partial_k\partial_l\sigma(0),
 \end{aligned}$$

where Δ_{n-1} is the Laplacian in Euclidean coordinates of \mathbb{R}^{n-1} (see [12, Section 2] for more details). We will now specify one more property of σ . We choose $r > 0$ small and we require that $\sigma(s) = |s|^4$ for $s \in U_r^\pm = \{s \in U^\pm : |s| < r\}$; if r is small enough then this is compatible with the fact that $\sigma \in h^{4+\alpha}(\Sigma) \cap W$. We conclude that

$$G(\sigma)|_{U^\pm}(0) = -24 < 0. \tag{2.3}$$

It follows from Proposition 2.1 that the evolution equation (2.2) with initial value $\rho(0) = \sigma$ has a unique solution

$$\rho(\cdot, \sigma) \in C([0, T_0], h^{4+\alpha}(\Sigma)) \cap C^1([0, T_0], h^\alpha(\Sigma)). \tag{2.4}$$

Next we consider the restriction $\rho^\pm(t, \sigma)$ on U^\pm of the function $\rho(t, \sigma)$, that is, $\rho^\pm(t, \sigma) := \rho(t, \sigma)|_{U^\pm}$ for $0 \leq t \leq T_0$, and we set $d^\pm(t) := \rho^\pm(t, \sigma)(0)$, to track the position of the center. It follows from (2.4) that $d^\pm \in C^1([0, T_0])$. Moreover, using the local character of G , we conclude that d^\pm satisfies the equation

$$(d^\pm)'(t) = G(\rho(t, \sigma))|_{U^\pm}(0) \quad \text{for } 0 \leq t \leq T_0, \quad d^\pm(0) = 0. \tag{2.5}$$

Equations (2.3)–(2.5) and the mean value theorem yield

$$d^\pm(t) = -Mt + \left(\int_0^1 ((d^\pm)'(\tau t) - (d^\pm)'(0)) d\tau \right) t, \tag{2.6}$$

where $M := 24$. It follows from (2.6) that there exists a positive constant $\mu > 0$ and an interval $(t_0, t_1) \subset (0, T_0]$ such that $\rho^\pm(t, \sigma)(0) = d^\pm(t) \leq -\mu$ for $t \in (t_0, t_1)$. By Proposition 2.1(b) we can find a function $\sigma_0 \in W_0$ such that $\Sigma_0 := \Gamma_{\sigma_0}$ is embedded and such that $\Gamma(t) := \Gamma_{\rho(t, \sigma_0)}$ is immersed for at least $t \in (t_0, t_1)$. By employing Proposition 2.1(b) once more we conclude

there is a neighborhood $W(\sigma_0) \subset W_0$ of σ_0 in $h^{2+\beta_0}(\Sigma)$ such that Γ_{ρ_0} is still embedded, whereas $\Gamma_{\rho(t,\rho_0)}$ is immersed for $t \in (t_0, t_1)$ and all $\rho_0 \in W(\sigma_0)$. We note that $C^{2+\beta}(\Sigma)$ is contained in $h^{2+\beta_0}(\Sigma)$ with continuous injection $j : C^{2+\beta}(\Sigma) \rightarrow h^{2+\beta_0}(\Sigma)$. Hence $U_0 := j^{-1}(W(\sigma_0))$ is an open $C^{2+\beta}$ -neighborhood of σ_0 , and Theorem 1 follows by choosing $\Sigma_0 := \Gamma_{\sigma_0}$ and $\Gamma_0 := \Gamma_{\rho_0}$ for $\rho_0 \in U_0$. \square




Figure 2 This is half of Γ_0 , a surface that loses embeddedness and becomes immersed. The gap might have to be much smaller than depicted.

3. The volume-preserving mean curvature flow. As in the previous section Σ denotes a smooth, compact, closed, immersed, orientable hypersurface in \mathbb{R}^n , and we define $W := \{\rho \in h^{1+\beta_0}(\Sigma) : \|\rho\|_\infty < a\}$ for $a > 0$ appropriate. The volume-preserving mean curvature flow (1.2) in \mathcal{R} is equivalent to the following evolution equation for the distance function ρ :

$$\partial_t \rho = G(\rho), \quad \rho(0) = \rho_0, \tag{3.1}$$

where $\rho_0 \in W$ is chosen such that $\Gamma_0 = \Gamma_{\rho_0}$, and where

$$G(\rho) := L(\rho)(\overline{H}_{\Gamma_\rho} - \theta_\rho^* H_{\Gamma_\rho}), \quad \rho \in h^{2+\alpha}(\Sigma) \cap W. \tag{3.2}$$

Here H_{Γ_ρ} is the mean curvature of Γ_ρ and $L(\rho)$ comes from calculating the normal velocity in coordinates of Σ ; see [14] for more details. We have the following existence result for solutions of (3.1).

Proposition 3.1.

(a) *Let $\sigma \in W$ be given. Then there exists a positive constant $T_0 > 0$ and a neighborhood $W_0 \subset W$ of σ in $h^{1+\beta_0}(\Sigma)$ such that (3.1) has a unique solution,*

$$\rho(\cdot, \rho_0) \in C([0, T_0], W) \cap C^\infty((0, T_0) \times \Sigma) \text{ for every } \rho_0 \in W_0.$$

- (b) The map $[(t, \rho_0) \mapsto \rho(t, \rho_0)]$ defines a smooth, local semiflow on W_0 .
- (c) $\rho(\cdot, \rho_0) \in C([0, T_0], h^{2+\alpha}(\Sigma)) \cap C^1([0, T_0], h^\alpha(\Sigma))$ for every $\rho_0 \in h^{2+\alpha}(\Sigma) \cap W_0$.

Proof. Part (a) and part (b) follow from the results in [14, Section 2]. To be more precise, in [14] only embedded surfaces are considered. However, a careful analysis of the proof shows that the existence, uniqueness, and semiflow results remain valid for immersed hypersurfaces, provided one defines the mappings X and Φ_ρ of [14, Section 2] by their local analogs as in [12, Section 2]. Part (c) can be established by similar arguments as in the proof of Proposition 2.1. \square

We proceed to prove Theorem 2. Our first goal is to construct a suitable reference manifold Σ . We take a positively oriented, immersed curve in \mathbb{R}^2 , such as the one in Figure 3. The immersed image contains a line segment twice, with opposite orientations, and the image without this line segment is an embedded curve.




Figure 3 Rotation of this curve in \mathbb{R}^n yields the hypersurface Σ .

For the two-dimensional case this curve will be Σ , while for the higher-dimensional case we rotate the curve to generate a hypersurface, as outlined below. Let $[s \mapsto (x(s), y(s))] : [0, L] \rightarrow \mathbb{R}^2$ be a parametrization by arc length of the curve and let $S^{n-2} \subset \mathbb{R}^{n-1}$ be the standard $(n - 2)$ -dimensional unit sphere. Then we set

$$\Sigma = \{(x(s), y(s)\omega) : s \in [0, L], \omega \in S^{n-2}\}.$$

Let κ_1 denote the scalar curvature of the curve $[s \mapsto (x(s), y(s))]$. Then a standard computation yields the mean curvature of Σ as

$$H = \frac{1}{n-1} \left(\kappa_1 - (n-2) \frac{x'}{y} \right). \tag{3.3}$$

Furthermore, the symmetry of Σ can be used to compute the average of H as

$$\overline{H} = \frac{|S^{n-2}|}{|\Gamma|} \int_0^L H(s)y(s) ds.$$

Using equation (3.3) and setting $d = \min\{y(s)\}$ one derives

$$\begin{aligned} (n-1) \int_0^L H(s)y(s) ds &= d \int_0^L \kappa_1(s) ds + \int_0^L \kappa_1(s)(y(s) - d) ds \\ &\quad - (n-2) \int_0^L x'(s) ds. \end{aligned}$$

The theorem of the turning tangents implies that $\int \kappa_1(s) ds = 2\pi$, and hence it is clear that $\overline{H} > 0$ provided d is large enough, which amounts to shifting the curve far enough away from the axis of rotation. By continuity the average of the mean curvature of Γ_ρ is therefore also positive provided $\rho \in W$ is small enough. Finally, it is clear that by construction Σ contains a flat $(n-1)$ -dimensional annulus U twice, with opposite orientations, and that Γ_ρ is in fact embedded in \mathbb{R}^n provided $\rho < 0$ on the annulus.

We let U^\pm be the two components of $i^{-1}(U)$, where $i : \Sigma \rightarrow \mathbb{R}^n$ is the immersion under consideration, and U is the flat annulus from above. We now choose $\sigma \in h^{2+\alpha}(\Sigma) \cap W$ with $\sigma \equiv 0$ on U^\pm (in fact we could choose $\sigma \equiv 0$ on Σ so that $\Gamma_\sigma = \Sigma$); then by (3.2)

$$G(\sigma)|_{U^\pm} = \overline{H}_{\Gamma_\sigma} > 0. \tag{3.4}$$

Let $\rho(\cdot, \sigma)$ be the unique solution of (3.1) with initial value $\rho(0) = \sigma$, and note that

$$\rho(\cdot, \sigma) \in C([0, T_0], h^{2+\alpha}(\Sigma)) \cap C^1([0, T_0], h^\alpha(\Sigma)) \tag{3.5}$$

due to Proposition 3.1. As in Section 2 we let $\rho^\pm(t, \sigma)$ denote the restriction of $\rho(t, \sigma)$ on U^\pm . We set $d^\pm(t) := \rho^\pm(t, \sigma)(s_0, \omega)$ with s_0 any fixed point on

the line segment that generated U and any fixed $\omega \in S^{n-2}$. It follows that $d^\pm \in C^1[0, T_0]$, and it is easy to see that d^\pm solves the equation

$$(d^\pm)'(t) = G(\rho(t, \sigma))|_{U^\pm}(s_0, \omega) \quad \text{for } 0 \leq t \leq T_0, \quad d^\pm(0) = 0. \quad (3.6)$$

Equations (3.4)–(3.6) and the mean value theorem show that

$$d^\pm(t) = Mt + \left(\int_0^1 ((d^\pm)'(\tau t) - (d^\pm)'(0)) d\tau \right) t$$

with $M := \overline{H}_{\Gamma_\sigma}$. Using Proposition 3.1(b) we can choose a function σ_0 in W_0 such that Γ_{σ_0} is embedded and such that $\Gamma_{\rho(t, \sigma_0)}$ ceases to be embedded on a time interval $(t_0, t_1) \subset (0, T_0]$. The idea is that the time derivative of d^\pm is positive, and hence so will be d^\pm for some later time even if it was initially negative; see Section 2 for more details. According to Proposition 3.1(b) the same behavior will still prevail for ρ_0 in a small enough $h^{1+\beta_0}(\Sigma)$ -neighborhood $W(\sigma_0) \subset W_0$ of σ_0 . Theorem 2 now follows by setting $U_0 := j^{-1}(W(\sigma_0))$ with $j := C^{1+\beta}(\Sigma) \rightarrow h^{1+\beta_0}(\Sigma)$ the natural injection.

Acknowledgment. The research of the second author has been partially supported by NSF, Grant DMS-9801337.

REFERENCES

- [1] H. Amann, “Linear and Quasilinear Parabolic Problems,” Vol. I, Birkhäuser, Basel, 1995, Vol. II, III, in preparation.
- [2] S.B. Angenent, *Nonlinear analytic semiflows*, Proc. Roy. Soc. Edinburgh Sect. A **115** (1990), 91–107.
- [3] A.J. Bernoff, A.L. Bertozzi, and T.P. Witelski, *Axisymmetric surface diffusion: Dynamics and stability of self-similar pinch-off*, J. Statist. Phys., **93** (1998), 725–776.
- [4] P. Baras, J. Duchon, and R. Robert, *Évolution d’une interface par diffusion de surface*, Comm. Partial Differential Equations **9** (1984), no. 4, 313–335.
- [5] L. Bronsard and B. Stoth, *Volume-preserving mean curvature flow as a limit of a nonlocal Ginzburg-Landau equation*, SIAM J. Math. Anal. **28** (1997), 769–807.
- [6] J.W. Cahn, C.M. Elliott, and A. Novick-Cohen, *The Cahn–Hilliard equation with a concentration dependent mobility: motion by minus the Laplacian of the mean curvature*, European J. Appl. Math. **7** (1996), 287–301.
- [7] J.W. Cahn and J.E. Taylor, *Surface motion by surface diffusion*, Acta metall. mater. **42** (1994), 1045–1063.

- [8] B.D. Coleman, R.S. Falk, and M. Moakher, *Space-time finite element methods for surface diffusion with applications to the theory of the stability of cylinders*, SIAM J. Sci. Comput. **17** (1996), 1434–1448.
- [9] G. DaPrato and P. Grisvard, *Equations d'évolution abstraites nonlinéaires de type parabolique*, Ann. Mat. Pura Appl. (4) **120** (1979), 329–396.
- [10] F. Davì and M.E. Gurtin, *On the motion of a phase interface by surface diffusion*, Z. Angew. Math. Phys. **41** (1990), 782–811.
- [11] C.M. Elliott and H. Garcke, *Existence results for geometric interface models for surface diffusion*, Adv. Math. Sci. Appl. **7** (1997), 467–490.
- [12] J. Escher, U.F. Mayer, and G. Simonett, *The surface diffusion flow for immersed hypersurfaces*, SIAM J. Math. Anal. **29** (1998), 1419–1433.
- [13] J. Escher, U.F. Mayer, and G. Simonett, *On the surface diffusion flow*, Proc. Intern. Conf. on Navier–Stokes Equations and Related Problems, TEV/VSP, Vilnius/Utrecht, 1998.
- [14] J. Escher and G. Simonett, *The volume preserving mean curvature flow near spheres*, Proc. Amer. Math. Soc. **126** (1998), 2789–2796.
- [15] M. Gage, *On an area-preserving evolution equation for plane curves*, *Nonlinear Problems in Geometry*, D.M. DeTurck, editor, Contemp. Math. **51**, AMS, Providence (1986), 51–62.
- [16] Y. Giga and K. Ito, *On pinching of curves moved by surface diffusion*, Comm. Appl. Anal. **2** (1998), 393–405.
- [17] G. Huisken, *The volume preserving mean curvature flow*, J. Reine Angew. Math. **382** (1987), 35–48.
- [18] A. Lunardi, “Analytic semigroups and optimal regularity in parabolic problems,” Birkhäuser, Basel, 1995.
- [19] U.F. Mayer, *Numerical simulations for the surface diffusion flow in three space dimensions*, preprint.
- [20] W.W. Mullins, *Theory of thermal grooving*, J. Appl. Phys. **28** (1957), 333–339.
- [21] A. Polden, *Curves and surfaces of least total curvature and fourth-order flows*, Ph.D. dissertation, Universität Tübingen, Germany (1996).